arXiv:1112.4181v1 [math.DG] 18 Dec 2011

Einstein Hermitian Metrics of Positive Sectional Curvature

Caner Koca

Abstract

In this paper we will prove that the only compact 4-manifold M with an Einstein metric of positive sectional curvature which is also hermitian with respect to some complex structure on M, is $\mathbb{CP}_2$, with its Fubini-Study metric.

1 Introduction

Let M be a smooth n-manifold. A Riemannian metric g on M is called *Einstein* if the Ricci tensor is a constant multiple of the metric tensor, i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$, called the *Einstein constant* [4]. If $\lambda > 0$ and g is complete, M is compact by Myers' Theorem. Since the Ricci tensor is, by definition, the pointwise average of all sectional curvatures, for an Einstein metric the positivity of λ is assured if all sectional curvatures are positive.

In this paper we are dealing with compact smooth manifolds in dimension 4 which admit Einstein metrics with $\lambda > 0$. Examples of such manifolds are 4-sphere S^4 with its standard round metric (which has all sectional curvatures $K \equiv 1$), and the complex projective plane $\mathbb{CP}_2$ with the Fubini-Study metric g_{FS} (which has $1 \leq K \leq 4$ everywhere). Notice that these examples are of strictly *positive* sectional curvature. The product metric on $S^2 \times S^2$ is Einstein with $\lambda > 0$, too, its sectional curvatures, however, are *non-negative*. They are actually 0 for transverse planes (i.e. $K_{\Pi} = 0$ if the plane Π is not tangent to each of the factors). In fact, the famous Hopf Conjecture asks whether or not there are any metrics on $S^2 \times S^2$ of positive sectional curvature.

The most fruitful resource of Einstein metrics is the Kähler geometry. For a compact complex surface M,

a. there is a unique Kähler-Einstein metric with $\lambda < 0$ if $c_1(M) < 0$ (see [22], [2]),

b. there is a unique Kähler-Einstein metric with $\lambda = 0$ (i.e. Ricci-flat) in each Kähler class if $c_1(M) = 0$ (see [21]).

c. In $\lambda > 0$ case, Tian [20] showed that M admits a Kähler-Einstein metric with $\lambda > 0$ iff M has $c_1(M) > 0$ and its automorphism group $\mathrm{Aut}(M)$ is a reductive Lie group. The diffeomorphism types of such complex surfaces are $\mathbb{CP}_2$, $\mathbb{CP}_1 \times \mathbb{CP}_1$ and $\mathbb{CP}_2 \sharp k \overline{\mathbb{CP}}_2$ with $k = 3, 4, \ldots, 8$.

Among those Kähler-Einstein metrics, only the ones on $\mathbb{CP}_2$ is of *positive* sectional curvature. This follows, for example, by Andreotti's Theorem [1], which says that any compact complex surface M with a Kähler metric of positive sectional curvature must be $\mathbb{CP}_2$. Andreotti's theorem is a special case of Frankel's conjecture [9], which was later proved by Siu and Yau [19], which asserts that the generalization of the statement is true for all complex dimensions n.

As a consequence of the following theorem, Gursky and LeBrun [10] reached at the same conclusion that $\mathbb{CP}_2$ is the unique compact complex surface with a positively curved Kähler-Einstein metric:

Theorem 1 (Gursky-LeBrun) *Let (M^4, g) be a compact oriented Einstein 4-manifold of non-negative sectional curvature.*

(i) if M has positive intersection form, then $(M, g) = (\mathbb{CP}_2, g_{FS})$, up to rescaling and isometry;

(ii) if g is neither self-dual nor anti-self-dual, then $\frac{15}{4}|\tau| < \chi \leq 9$ where τ is the signature and χ is the Euler characteristic of M.

Note that $\mathbb{CP}_2\sharp k\overline{\mathbb{CP}}_2$ have $\tau = 1-k$ and $\chi = 3+k$, so the inequality is not satisfied if $k = 3, 4, \ldots, 8$. The Kähler-Einstein metrics on those surfaces cannot be self-dual (that is, $W_- \equiv 0$) or anti-self-dual (that is, $W_+ \equiv 0$) as a consequence of

$$12\pi^2\tau = \int_M |W_+|^2 - |W_-|^2 d\mu \qquad \text{(Signature Formula)}$$

$$8\pi^2\chi = \int_M |W_+|^2 + |W_-^2| + \frac{s^2}{24} - \frac{|\mathring{r}|}{2} d\mu \qquad \text{(Gauss-Bonnet)}$$

and the fact that $|W_+|^2 = \frac{s^2}{24}$ for Kähler metrics. Indeed, since $\tau < 0$, the signature formula implies that g cannot be self-dual. If, on the other hand, g were anti-self-dual, then $s = 0$ since g is Kähler, and also $\mathring{r} = 0$ since g is Einstein. So, the two formulas would give different values for the integral $\int_M |W_-|^2 d\mu$ unless $k = 9$; but this case is excluded from the range for k.

Observe, in passing, that part (ii) has a similar taste with Hitchin's theorem [12], which says that a compact oriented Einstein 4-manifold of positive sectional curvature should satisfy $\chi \geq (\frac{3}{2})^{3/2}|\tau|$; and the Hitchin-Thorpe inequality $\chi \geq \frac{3}{2}|\tau|$ which holds for all compact orientable Einstein 4-manifolds.

If we relax the Kähler condition on the Einstein metric g, and merely assume that g is *hermitian*, that is $g(J\cdot, J\cdot) = g(\cdot,\cdot)$ for a complex structure J on the manifold M, interesting enough, we get only two more exceptional metrics:

Theorem 2 (LeBrun [15]) *Let (M^4, J) be a compact complex surface. If g is Einstein and Hermitian, then only one of the following holds:*

(1) g is Kähler-Einstein with $\lambda > 0$.

(2) (M, J) is biholomorphic to $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$ and g is the Page metric g_{Page} (up to rescaling and isometry).

(3) (M, J) is biholomorphic to $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ and g is the Chen-LeBrun-Weber metric g_{CLW} (up to rescaling and isometry).

Thus, if we in addition assume that g is of positive sectional curvature, then the first case of the above theorem is possible only when M is $\mathbb{CP}_2$, by Andreotti's theorem. Moreover, by a theorem of Berger [3], the Kähler-Einstein metric g on $\mathbb{CP}_2$, is the Fubini-Study metric (up to rescaling and isometry) since it has positive holomorphic bisectional curvature.

In the next two sections we will prove that the two exceptional metrics g_{Page} and g_{CLW} are not of positive sectional curvature either. This will conclude the proof of our main theorem:

Theorem 3 *Let M^4 be a compact smooth 4-manifold, and let g be an Einstein metric of positive sectional curvature. If g is hermitian with respect to some complex structure J on M, then (M, J) is biholomorphic to $\mathbb{CP}_2$, and g is the Fubini-Study metric (up to rescaling and isometry).*

One of the key facts in the proof of this theorem is Frankel's Theorem [9] which says that totally geodesic submanifolds of complementary dimensions on positively curved manifolds necessarily intersect. Since the Page metric has an explicit form, we are also able to give a computational proof of the failure of positivity. Note that, on the contrary, Chen-LeBrun-Weber metric does not possess such an explicit formula.

2 Page metric

The Page metric was first introduced by D. Page in 1978 as a limiting metric of Kerr-de Sitter solution (see [17]). To define it formally, we first think of the following metric on the product $S^3 \times I$ where I is the closed interval $[0, \pi]$:

$$g = V(r)dr^2 + f(r)(\sigma_1^2 + \sigma_2^2) + \frac{C\sin^2 r}{V(r)}\sigma_3^2$$

where the coefficient functions are given as

$$V(r) = \frac{1 - a^2 \cos^2 r}{3 - a^2 - a^2(1 + a^2) \cos^2 r}$$

$$f(r) = \frac{4}{3 + 6a^2 - a^4}(1 - a^2 cos^2 r)$$

$$C = \left(\frac{2}{3 + a^2} \right)^2$$

and a is the unique positive root of $a^4 + 4a^3 - 6a^2 + 12a - 3 = 0$. Here, $\sigma_1, \sigma_2, \sigma_3$ is the standard left invariant 1-forms on the Lie group $SU(2) \approx S^3$.

When $r = 0$ or π, we see from the formula that the metric reduces to a round metric on S^2. Thus, g descends to a metric, denoted by g_{Page}, on the quotient $(S^3 \times I)/ \sim$ where $\sim$ identifies the fibers of the Hopf fibration $p : S^3 \to S^2$ on the two ends $S^3 \times \{0\}$ and $S^3 \times \{\pi\}$ of the cylinder $S^3 \times I$.



The resulting manifold is indeed the connected sum $\mathbb{CP}_2 \sharp \overline{\mathbb{CP}}_2$. To see this, recall that in the cell decomposition of $\mathbb{CP}_2$, the attaching map from the boundary of the 4-cell (which is S^3) to the 2-skeleton (which is $\mathbb{CP}_1 \approx S^2$) is given by the Hopf map [11]. So, if we cut the cylinder $S^3 \times I$ in two halves and identify the Hopf fibers of S^3 at each end, we get $\mathbb{CP}_2 - \{\text{small ball}\}$. Since the right and left halves have different orientations, we obtain $\mathbb{CP}_2 \sharp \overline{\mathbb{CP}}_2$ in the quotient.



Now, we will prove that the Page metric is not of positive sectional curvature. We will use the following classical theorem by Frankel:

Theorem 4 (Frankel [9]) *Let M be a smooth n-manifold, and let g be a complete Riemannian metric of positive sectional curvature. If X and Y are two compact totally geodesic submanifolds of dimensions d_1 and d_2 such that $d_1 + d_2 \geq n$, then X and Y intersect.*

In our case, the two 2-spheres on each end of the above quotient will play the role of X and Y. They are compact and the dimensions add up to 4. So it remains to show that those two submanifolds are totally geodesic with respect to g_{Page}. Since they are obviously disjoint, this will imply that g_{Page} cannot have positive sectional curvature.

There is a very well-known lemma to detect totally geodesic submanifolds:

Lemma 1 *Let (M, g) be a Riemannian manifold. If f is an isometry, then each connected component of the fix point set* $\mathrm{Fix}(f)$ *of f is a totally geodesic submanifold of M.*

So, below we will show that there is an isometry of the Page metric whose fix point set is precisely the two end spheres.

What are the isometries of the Page metric? Derdziński [8] showed that the Page metric is indeed conformal to one of Calabi's extremal

Kähler metrics on $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$. On the other hand, the identity component of the isometry group of extremal Kähler metrics is a maximal compact subgroup of the identity component of the automorphism group [6]. In the case of $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$, this implies that the identity component of the isometry group of the Page metric is $U(2) = (SU(2) \times S^1)/\mathbb{Z}_2$. By the formula of the metric, we see that the isometries in the $SU(2)$ component are precisely given by the left multiplication action of $SU(2)$ on the first factor of $S^3 \times I$. Note that the forms σ_i, $i = 1, 2, 3$ are invariant under the action, but the action on the 3-spheres $S^3 \times \{r\}$, $r \in (0, \pi)$ is fixed-point-free! The metric is invariant under this action as the coefficients of the metric only depend on the parameter r.

Now, let us see what happens at the endpoints $r = 0$ and $r = \pi$: It is well-known that the action of $U \in SU(2)$ on the 2-sphere S^2 (after the quotient) is given by the conjugation $A \mapsto UAU^{-1}$, where we regard the 2×2 complex matrix $A = x\sigma_1 + y\sigma_2 + z\sigma_3$ with $x^2 + y^2 + z^2 = 1$ as a point of S^2. It is now straightforward to see that the action of $-I \in SU(2)$ is trivial on S^2 (since $(-I)A(-I)^{-1} = A$); thus, it fixes every point on S^2. Therefore, we conclude that the fixed point set of the isometry given by the "antipodal map" $-I \in SU(2)$ consists of the two 2-spheres at each end of the quotient $\big((S^3 \times I)/\sim\big) \approx \mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$. Note that, indeed, there is an S^1-family of isometries generated by rotation in direction of σ_3 having the exact same fixed point set.

So we showed that there are two disjoint compact totally geodesic submanifolds of $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$. Therefore, by Frankel's theorem, we conclude that g_{Page} is not of positive sectional curvature.

Finally, we note that we can actually show the failure of positivity *directly by brute-force* using tensor calculus: Introduce a new coordinate function $x := \cos(r)$, so that the metric becomes

$$g = W^2(x)dx^2 + g^2(x)(\sigma_1^2 + \sigma_2^2) + \frac{D^2}{W(x)}\sigma_3^2$$

where the coefficient functions are given as

$$
\begin{aligned}
W(x) &= \sqrt{\frac{1-a^2x^2}{(3-a^2-a^2(1+a^2)x^2)(1-x^2)}}\\
g(x) &= 2\sqrt{\frac{1-a^2x^2}{3+6a^2-a^4}}\\
D &= \frac{2}{3+a^2}
\end{aligned}
$$

and choose the following vierbein: $\{Wdx, g\sigma_1, g\sigma_2, \frac{D}{W}\sigma_3\} =: \{e^0, e^1, e^2, e^3\}$. Then by a standard tensor calculus, we see that the sectional curvature of the plane generated by e_0 and e_1 is given by

$$K_{01} = 2\frac{g'W' - g''W}{gW^3}.$$

Using a computer program like *Maple*, one can easily verify that this function $K_{01}(x)$ can take both positive and negative values for $x \in (-1, 1)$.

3 Chen-LeBrun-Weber metric

After the discovery of the Kähler-Einstein metrics on $\mathbb{CP}_2\sharp k\overline{\mathbb{CP}}_2$ for $k = 3, \ldots, 8$, and the Einstein metric (namely the Page metric) on $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$, it was speculated that whether $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ admits an Einstein metric. Derdziński [8] had discovered in early '80s that even though Page metric is not Kähler, it is actually confomally related to a Kähler metric on $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$; indeed to one of Calabi's extremal Kähler metrics in [5]. Inspired by this result, LeBrun showed in 1995 that an Einstein hermitian metric h on $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ has to be conformally related to an extremal Kähler metric [14], in such a way that $h = s^{-2}g$, where s is the scalar curvature of g, which turns out to be necessarily positive in this setting. Conversely, it was proved that for an extremal Kähler metric g, the metric $h := s^{-2}g$ is Einstein (defined wherever $s \neq 0$) if g is the critical point of the Calabi functional regarded as an action on extremal Kähler cone [18], [7].

Unlike the $\mathbb{CP}_2\sharp\overline{\mathbb{CP}}_2$ case, not every Kähler class on $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ is represented by an extremal Kähler metric. Nevertheless, using the computations of Futaki invariant in [13], Chen, LeBrun and Weber [7] showed that this action has a critical point, and this critical class is indeed represented by an *extremal* Kähler metric g of positive scalar curvature! Thus, the conformally related metric $s^{-2}g$ is an Einstein metric on $\mathbb{CP}_2\sharp\overline{2}\mathbb{CP}_2$, denoted by g_{CLW}. More recenty, LeBrun [15] showed that this is the *unique* Einstein hermitian metric on $\mathbb{CP}_2\sharp\overline{2}\mathbb{CP}_2$, and it can be obtained also as an appropriate deformation of the Kähler-Einstein metric in $\mathbb{CP}_2\sharp 3\overline{\mathbb{CP}}_2$ representing the first Chern class c_1 (see [16]).

The identity component of the isometry group of g_{CLW} lies in the identity component of the group of biholomorphisms of $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ as a maximal compact subgroup. Indeed, there is a natural torus action [7] of $S^1 \times S^1$ on $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ obtained by lifting the torus action

$$(e^{i\theta}, e^{i\xi}) \longmapsto ([u_1 : e^{i\theta}u_2], [v_1 : e^{i\xi}v_2])$$

on $\mathbb{CP}_1 \times \mathbb{CP}_1$ to its blowup at $([0:1],[0:1])$. Note that $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$ is isomorphic to the blowup of $\mathbb{CP}_1 \times \mathbb{CP}_1$ at one point. This torus action has the following moment map profile [15]:



Here, x and y are Hamiltonians. In particular, the fix point set of the vector field of the Hamiltonian x contains the two vertical edges of the pentagon, which indeed correspond to two disjoint holomorphic $\mathbb{CP}_1$'s lying in $\mathbb{CP}_2\sharp 2\overline{\mathbb{CP}}_2$. Again, by Lemma 1, they are totally geodesic. Therefore, by Frankel's theorem, it follows that g_{CLW} is not of positive sectional curvature. This completes the proof of Theorem 3.

References

[1] Andreotti, A. *On the complex structures of a class of simply connected manifolds*, in the Lefschetz symposium volume Algebraic geometry and topology, Princeton, 1957.

[2] Aubin, T. *Nonlinear analysis on manifolds. Monge-Ampere equations.* Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 252. Springer-Verlag, New York, 1982.

[3] Berger, M. *Sur les varietes d'Einstein compactes*, Comptes Rendus de la IIIe Reunion du Groupement des Mathematiciens dExpression Latine (Namur, 1965) pp. 35-55.

[4] Besse, A., *Einstein Manifolds*, Springer-Verlag, 1987.

[5] Calabi, E. *Extremal Kähler metrics*. Seminar on Differential Geometry, pp. 259-290, Ann. of Math. Stud., 102, Princeton Univ. Press, Princeton, N.J., 1982.

[6] Calabi E. *Extremal Kähler metrics II*, in Differential Geometry and Complex Analysis, Springer, Berlin, 1985, pp. 95-114.

[7] Chen, X.X., LeBrun, C., Weber, B. *On Conformally Kähler, Einstein Manifolds*, J. Amer. Math. Soc. 21 (2008), no. 4, 1137-1168.

[8] Derdziński A. *Self-dual Kähler manifolds and Einstein manifolds of dimension four*, Compositio Math., 49 (1983), pp. 405-433.

[9] Frankel, T. *Manifolds with positive curvature*. Pacific J. Math. 11 1961 165-174.

[10] Gursky, M.J., LeBrun, C. *On Einstein manifolds of positive sectional curvature*. Ann. Global Anal. Geom. 17 (1999), no. 4, 315-328.

[11] Hatcher, A. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.

[12] Hitchin N.J., *On Compact Four-Dimensional Einstein Manifolds*, J. Diff. Geom. 9 (1974) 435-442.

[13] LeBrun, C., Simanca, S. R. *Extremal Kähler metrics and complex deformation theory*. Geom. Funct. Anal. 4 (1994), no. 3, 298-336.

[14] LeBrun, C. *Einstein metrics on complex surfaces*. Geometry and physics (Aarhus, 1995), 167-176, Lecture Notes in Pure and Appl. Math., 184, Dekker, New York, 1997.

[15] LeBrun, C. *On Einstein, Hermitian 4-Manifolds*, e-print arXiv:1010.0238 [math.DG].

[16] LeBrun, C. *Einstein Manifolds and Extremal Kähler Metrics*, e-print arXiv:1009.1270 [math.DG]

[17] Page, D. *A Compact Rotating Gravitational Instanton*, Physics Letters, Vol. 79B, no.3, (1978), 235–238.

[18] Simanca, S.R. *Strongly extremal Kähler metrics*, Ann. Global Anal. Geom., 18 (2000), pp. 29-46.

[19] Siu, Y.-T., Yau, S.-T., *Compact Kähler manifolds of positive bisectional curvature*, Invent. Math. 59 (1980), 189-204.

[20] Tian, G. *On Calabi's conjecture for complex surfaces with positive first Chern class.* Invent. Math. 101 (1990), no. 1, 101-172.

[21] Yau, S.-T. *Calabi's conjecture and some new results in algebraic geometry*. Proc. Nat. Acad. Sci. U.S.A. 74 (1977), no. 5, 1798-1799.

[22] Yau, S.-T. *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampre equation.* I. Comm. Pure Appl. Math. 31 (1978), no. 3, 339-411.

Acknowledgement. I would like to thank my advisor Claude LeBrun for suggesting the problem and for his help, guidance and encouragement. Also many thanks to Selin Taşkent, Yongsheng Zhang and Mustafa Kalafat for useful discussions and comments.

AUTHOR'S ADDRESS:

Mathematics Department, SUNY, Stony Brook, NY 11794, USA

AUTHOR'S E-MAIL:

`caner@math.sunysb.edu`